## **Guidance for HCW PPE During COVID-19 Response**

- Negative pressure rooms are not required as the standard care environment for patients with known or suspected COVID-19.
- Aerosol-generating procedures (e.g., bronchoscopy, cardiopulmonary resuscitation, endotracheal intubation and extubation, manual ventilation, non-invasive ventilation such as BiPAP or CPAP, open suctioning of airways, sputum induction) for patients with suspected/confirmed COVID-19 should be performed in negative pressure rooms or rooms with HEPA filter when available. When unavailable, all healthcare workers entering the room must wear respiratory protection for the recommended time frame after the aerosol-generating procedure.
- Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient
- **NOTE:** These are INTERIM RECOMMENDATIONS and are expected to change. Recommended PPE reflects the minimum recommended PPE. Use additional PPE per Standard and Transmission-Based Precautions Policy.

Inpatient				
Patient Scenario	HCW Provider type	Recommended PPE	Preferred type of room	
Patient with confirmed/suspected COVID-19 AND ICU or tenuous respiratory status	All in room  Minimize number of people entering room	<ul> <li>Special Airborne Precautions</li> <li>N95/PAPR</li> <li>Eye protection</li> <li>Gown</li> <li>Gloves</li> <li>Shoe covers</li> <li>Patient face mask*</li> </ul>	<ul> <li>Negative pressure or HEPA filter, if available</li> <li>If non-negative pressure and NO HEPA filter, then healthcare workers should wear N95/PAPR when entering the room of patient for 2 hours after aerosol-generating procedures</li> </ul>	
Patient with confirmed/suspected COVID-19 AND Non-ICU	All in room  Minimize number of people entering room	Special Airborne Precautions  N95/PAPR  Eye protection  Gown  Gloves  Shoe covers  Patient face mask*	<ul> <li>Regular room</li> <li>Room does not need to remain closed after use</li> </ul>	
Patient without symptoms concerning for COVID-19  OR  Negative periprocedural test	All in room	Standard transmission-based precautions  • Face mask • Patient face mask*	<ul> <li>Regular room</li> <li>Room does not need to remain closed after use</li> </ul>	

<sup>\*</sup>Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient.



Emergency Department				
Patient Scenario	HCW Provider type	Recommended PPE	Preferred type of room	
Patient with confirmed/suspected COVID-19  AND Tenuous respiratory status	All	<ul><li>Follow Inpatient Guidelines</li><li>Patient face mask*</li></ul>	Follow Inpatient Guidelines	
Patient with pending respiratory failure requiring intubation	All	Special Airborne Precautions  N95/PAPR  Eye protection  Gown  Gloves  Patient face mask*	<ul> <li>Negative pressure or HEPA filter, if available</li> <li>If non-negative pressure or NO HEPA filter, then healthcare workers should wear N95/PAPR when entering the room of patient for 2 hours after aerosol-generating procedures</li> </ul>	
	Triage (if initiating treatment follow RN/MD)	<ul><li>Face mask</li><li>Patient face mask*</li></ul>		
Patient with fever or cough/shortness of breath concerning for COVID-19	HCW performing assessment/providing treatment	<ul> <li>Face mask</li> <li>Gown</li> <li>Gloves</li> <li>Patient face mask*</li> </ul>	<ul> <li>Single room with door closed</li> <li>Room does not need to be closed after use</li> </ul>	
	HCW collecting nasal or respiratory specimens or performing nebulizer treatments	<ul> <li>N95/PAPR</li> <li>Eye protection</li> <li>Gown</li> <li>Gloves</li> <li>Patient face mask*</li> </ul>		
Patient without symptoms concerning for COVID-19	All	Standard transmission-based precautions  • Face mask  • Patient face mask*	<ul> <li>Regular room</li> <li>Room does not need to remain closed after use</li> </ul>	

<sup>\*</sup>Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient.



Procedures**				
Patient Scenario	HCW Provider type	Recommended PPE	Preferred type of room	
Patient with confirmed/or unknown COVID-19 status undergoing procedure (as defined below)	All in room  Minimize number of people entering room	<ul> <li>Special Airborne Precautions</li> <li>N95/PAPR</li> <li>Eye protection</li> <li>Gown</li> <li>Gloves</li> <li>Shoe covers</li> <li>Patient face mask*</li> </ul>	Per area-specific COVID plan	
Patient with negative peri- procedural COVID-19 test undergoing procedure	All in room	Standard transmission-based precautions      Face mask     Eye protection     Gown     Gloves     Patient face mask*	<ul> <li>Regular room</li> <li>Room does not need to remain closed after use</li> </ul>	

<sup>\*</sup>Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient.



<sup>\*\*</sup>These PPE recommendations apply to the following procedures: bronchoscopy, transesophageal echocardiography, electrical cardioversion, electrophysiology procedures requiring general anesthesia, electroconvulsive therapy, upper and lower endoscopy, fluoroscopically guided enteric tube placements, and interventional radiology procedures requiring anesthesia or in patients with a tracheostomy. For patients with positive tests, the provider should carefully review indications for the procedure and consider deferring the procedure if possible. If the procedure is necessary, the procedure should proceed with N95, eye protection, gown, glove, and shoe covers. If testing is not available at the time of the procedure, the procedure may proceed with N95, eye protection, gown, glove, and shoe covers.

Outpatient Respiratory Care Clinic			
Patient Scenario	HCW Provider type	Recommended PPE	Preferred type of room
Patient with fever or cough/shortness of breath concerning for COVID-19	Triage	Face mask	
		<ul> <li>Patient face mask*</li> </ul>	
	HCW performing assessment	N95/PAPR	
		Eye protection	
		• Gown	
		• Gloves	<ul> <li>Single room with door closed</li> </ul>
		<ul> <li>Patient face mask*</li> </ul>	<ul> <li>Room does not need to be closed after use</li> </ul>
	HCW collecting nasal or	• N95/PAPR	
	respiratory specimens or	Eye protection	
	administering nebulizer	• Gown	
	treatments	• Gloves	
		<ul> <li>Patient face mask*</li> </ul>	
		Standard transmission-based	
Patient without symptoms concerning for COVID-19	A.II	precautions	Regular room
	All	Face mask	Room does not need to remain closed after use
		<ul> <li>Patient face mask*</li> </ul>	

<sup>\*</sup>Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient.



Outpatient Non-Respiratory Care Clinic/Urgent Care			
Patient Scenario	HCW Provider type	Recommended PPE	Preferred type of room
	Triage	Face mask     Patient face mask*	
Patient with fever or cough/shortness of breath concerning for COVID-19	HCW performing assessment  HCW collecting nasal or respiratory specimens or administering nebulizer treatments	<ul> <li>Face mask</li> <li>Gown</li> <li>Gloves</li> <li>Patient face mask*</li> <li>N95/PAPR</li> <li>Eye protection</li> <li>Gown</li> <li>Gloves</li> <li>Patient face mask*</li> </ul>	Single room with door closed     Room does not need to be closed after use
Patient without symptoms concerning for COVID-19	All	Standard transmission-based precautions  • Face mask  • Patient face mask*	<ul> <li>Regular room</li> <li>Room does not need to remain closed after use</li> </ul>

<sup>\*</sup>Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient.



Outpatient/Inpatient Radiology Departments			
Patient Scenario	HCW Provider type	Recommended PPE	Preferred type of room
Patient with suspected/confirmed COVID-19	Radiology Provider	<ul> <li>Face mask</li> <li>Gown</li> <li>Gloves</li> <li>Patient face mask*</li> <li>Technicians performing portable x-rays in rooms of known/suspected patients follow special airborne precautions</li> </ul>	<ul> <li>Any radiology room location</li> <li>Room does not need to be closed after use</li> </ul>
Patient without symptoms concerning for COVID-19	All	Standard transmission-based precautions  • Face mask  • Patient face mask*	<ul> <li>Regular room</li> <li>Room does not need to remain closed after use</li> </ul>

6



<sup>\*</sup>Patient face masks should be worn by patients when they are outside of closed rooms as long as tolerated by the patient.